

Composition-Diamond Lemma for Differential Algebras*

Yuqun Chen, Yongshan Chen and Yu Li

School of Mathematical Sciences

South China Normal University

Guangzhou 510631

P. R. China

yqchen@scnu.edu.cn

jackalshan@126.com

liyu820615@126.com

Abstract: In this paper, we establish the Composition-Diamond lemma for free differential algebras. As applications, we give Gröbner-Shirshov bases for free Lie-differential algebras and free commutative-differential algebras, respectively.

Key words: Gröbner-Shirshov basis, differential algebra, Lie algebra.

AMS Mathematics Subject Classification(2000): 16S15, 13P10, 16S32

1 Introduction

In 1962, A. I. Shirshov [25] invented a theory that is now called the Gröbner-Shirshov bases theory for Lie algebras $Lie(X|S)$ presented by generators and defining relations. The main technical notion of Shirshov's theory was a notion of composition $(f, g)_w$ of two Lie polynomials, $f, g \in Lie(X)$ relative to some associative word w . Based on it, he defined an infinite algorithm of adding to some set S of Lie polynomials of all non-trivial compositions until one will get a set S^c that is closed under compositions, i.e., any non-trivial composition of two polynomials from S^c belongs to S^c (and leading associative words $\bar{s}$ of polynomials $s \in S^c$ do not contain each other as subwords). In addition, S and S^c generate the same ideal, i.e., $Id(S) = Id(S^c)$. S^c is now called a Gröbner-Shirshov basis of $Lie(X|S)$. Then he proved the following lemma.

Let $Lie(X) \subset \mathbb{k}\langle X \rangle$ be a free Lie algebra over a field $\mathbb{k}$ viewed as the algebra of Lie polynomials in the free algebra $\mathbb{k}\langle X \rangle$, and S a subset in $Lie(X)$. If $f \in Id(S)$, then $\bar{f} = u\bar{s}v$, where $s \in S^c$, $u, v \in X^$, $\bar{f}, \bar{s}$ are leading associative words of Lie polynomials f, s correspondingly, and X^* the free monoid generated by X .*

He used the following easy corollary of his lemma.

*Supported by the NNSF of China (No.10771077) and the NSF of Guangdong Province (No.06025062).

$Irr(S^c) = \{[u] \mid u \neq a\bar{s}b, s \in S^c, a, b \in X^*\}$ is a linear basis of the algebra $Lie(X|S) = Lie(X)/Id(S)$, where u is an associative Lyndon-Shirshov word in X^* and $[u]$ the corresponding non-associative Lyndon-Shirshov word under Lie brackets $[xy] = xy - yx$.

To define the Lie composition $(f, g)_w$ of two, say, monic Lie polynomials, where $\bar{f} = ac$, $\bar{g} = cb$, $c \neq 1$, a, b, c are associative words, and $w = acb$, A. I. Shirshov defines first the associative composition $fb - ag$. Then he puts on fb and ag special brackets $[fb]$, $[ag]$ in accordance with his paper [23]. The result is $(f, g)_w = [fc] - [cg]$. Following [25], one can easily get the same lemma for a free associative algebra: Let $S \subset \mathbb{k}\langle X \rangle$ and S^c be as before. If $f \in Id(S)$, then $\bar{f} = a\bar{s}b$ for some $s \in S^c$, $a, b \in X^*$. It was formulated later by L. A. Bokut [2] as an analogue of Shirshov's Lie composition lemma, and by G. Bergman [1] under the name "Diamond lemma" after the celebrated Newman's Diamond Lemma for graphs [22].

Shirshov's lemma is now called the Composition-Diamond lemma for Lie and associative algebras.

This kind of ideas were independently discovered by H. Hironaka [14] for power series algebras and by B. Buchberger [9, 10] for polynomial algebras. Buchberger suggested the name "Gröbner bases". Applications of Gröbner bases in mathematics (particularly in algebraic geometry) and in computer science are now well established.

At present, there are quite a few Composition-Diamond lemmas (CD-lemma for short) for different classes of non-commutative and non-associative algebras. Let us mention some.

A. I. Shirshov [24] proved a CD-lemma for commutative (anti-commutative) non-associative algebras, and mentioned that this lemma is also valid for non-associative algebras. It was used to solve the word problem for these classes of algebras. For non-associative algebras, this (but not the CD-lemma) was known, see A. I. Zhukov [27].

A. A. Mikhalev [19] proved a CD-lemma for Lie super-algebras.

T. Stokes [26] proved a CD-lemma for left ideals of an algebra $\mathbb{k}[X] \otimes E_{\mathbb{k}}(Y)$, the tensor product of an exterior (Grassman) algebra and a polynomial algebra.

A. A. Mikhalev and E. A. Vasilieva [20] proved a CD-lemma for the free supercommutative polynomial algebras.

A. A. Mikhalev and A. A. Zolotykh [21] proved a CD-lemma for $\mathbb{k}[X] \otimes \mathbb{k}\langle Y \rangle$, the tensor product of a polynomial algebra and a free algebra.

L. A. Bokut, Y. Fong and W. F. Ke [7] proved a CD-lemma for associative conformal algebras.

L. Hellström [15] proved a CD-lemma for a non-commutative power series algebra.

S.-J. Kang and K.-H. and Lee [16, 17] and E. S. Chibrikov [11] proved a CD-lemma for a module over an algebra (see also [12]).

D. R. Farkas, C. D. Feustel and E. L. Green [13] proved a CD-lemma for path algebras.

L. A. Bokut and K. P. Shum [8] proved a CD-lemma for Γ -algebras.

Y. Kobayashi [18] proved a CD-lemma for algebras based on well-ordered semigroups, L. A. Bokut, Yuqun Chen and Cihua Liu [5] proved a CD-lemma for dialgebras (see also [3]), L. A. Bokut, Yuqun Chen and Yongshan Chen [4] proved a CD-lemma for tensor product of free algebras, and L. A. Bokut, Yuqun Chen and Jianjun Qiu [6] proved a CD-lemma for associative algebras with multiple linear operators.

Let $\mathcal{D}$ be a set of symbols. Let $\mathbb{k}$ be a commutative ring with unit and $\mathcal{A}$ an associative algebra over $\mathbb{k}$. Then a pair $(\mathcal{A}, \partial_{\mathcal{A}})$ is called a differential algebra with differential operators $\mathcal{D}$ or $\mathcal{D}$ -algebra for short if $\partial_{\mathcal{A}} : \mathcal{D} \rightarrow \text{Der}\mathcal{A}$ is an injective map, where $\text{Der}\mathcal{A}$ stands for the set of all derivations of $\mathcal{A}$, i.e., any $\delta \in \text{Der}\mathcal{A}$, $a, b \in \mathcal{A}$,

$$\delta(ab) = \delta(a) \cdot b + a \cdot \delta(b).$$

Given a $\mathcal{D}$ -algebra $\mathcal{A}$ in this sense, we identify $D \in \mathcal{D}$ with the operator $\partial_{\mathcal{A}} D$.

In this paper, we establish a Composition-Diamond lemma for free differential algebras. As applications, we give Gröbner-Shirshov bases for free Lie-differential algebras and free commutative-differential algebras, respectively.

The authors would like to express their deepest gratitude to Professor L. A. Bokut for his kind guidance, useful discussions and enthusiastic encouragement. We also thank the referee for giving us some useful remarks.

2 Free differential algebras

In this section, we construct a free differential algebra with differential operators $\mathcal{D}$.

Let $\mathcal{A}$ be a $\mathcal{D}$ -algebra. A subset I of $\mathcal{A}$ is a $\mathcal{D}$ -ideal if I is a $\mathbb{k}$ -algebra ideal such that $D(I) \subseteq I$ for any $D \in \mathcal{D}$.

Let $\mathcal{A}, \mathcal{B}$ be $\mathcal{D}$ -algebras and $\psi : \mathcal{A} \rightarrow \mathcal{B}$ a map. Then ψ is called a $\mathcal{D}$ -homomorphism if ψ is a $\mathbb{k}$ -algebra homomorphism such that for any $a \in \mathcal{A}$ and $D \in \mathcal{D}$, $\psi(D(a)) = D(\psi(a))$, which is precisely $\psi(\partial_{\mathcal{A}} D(a)) = \partial_{\mathcal{B}} D(\psi(a))$.

Definition 2.1 *Let X be a set, $\mathcal{D}(X)$ a $\mathcal{D}$ -algebra and $\iota : X \rightarrow \mathcal{D}(X)$ an inclusion map. If for any $\mathcal{D}$ -algebra $\mathcal{A}$ and any map $\varphi : X \rightarrow \mathcal{A}$, there exists a unique $\mathcal{D}$ -homomorphism $\varphi^* : \mathcal{D}(X) \rightarrow \mathcal{A}$ such that $\varphi^* \iota = \varphi$, then $\mathcal{D}(X)$ is called a free $\mathcal{D}$ -algebra generated by X .*

Now we construct a free $\mathcal{D}$ -algebra directly.

Let $\mathcal{D} = \{D_j | j \in J\}$ and $\mathbb{N}$ the set of non-negative integers. For any $m \in \mathbb{N}$ and $\bar{j} = (j_1, \dots, j_m) \in J^m$, denote by $D^{\bar{j}} = D_{j_1} D_{j_2} \cdots D_{j_m}$ and $D^{\omega}(X) = \{D^{\bar{j}}(x) | x \in X, \bar{j} \in J^m, m \in \mathbb{N}\}$, where $D^{\bar{j}}(x) = x$ if $\bar{j} \in J^0$. Let $T = (D^{\omega}(X))^*$ be the free monoid generated by $D^{\omega}(X)$. For any $u = D^{\bar{i}_1}(x_1) D^{\bar{i}_2}(x_2) \cdots D^{\bar{i}_n}(x_n) \in T$, the length of u , denoted by $|u|$, is defined to be n . In particular, $|1| = 0$.

Let $\mathcal{D}(X) = \mathbb{k}T$ be the $\mathbb{k}$ -algebra spanned by T . For any $D_j \in \mathcal{D}$, we define the $\mathbb{k}$ -linear map $D_j : \mathcal{D}(X) \rightarrow \mathcal{D}(X)$ by induction on $|u|$ for $u \in T$:

- (1) $D_j(1) = 0$.
- (2) Suppose that $u = D^{\bar{i}}(x) = D_{i_1} D_{i_2} \cdots D_{i_{m_i}}(x)$. Then $D_j(u) = D_j D_{i_1} D_{i_2} \cdots D_{i_{m_i}}(x)$.
- (3) Suppose that $u = D^{\bar{i}}(x) \cdot v$, $v \in T$. Then $D_j(u) = (D_j D^{\bar{i}}(x)) \cdot v + D^{\bar{i}}(x) \cdot D_j(v)$.

By definition, D_j is a derivation on $\mathcal{D}(X)$ and then $\mathcal{D}(X)$ is a differential algebra with differential operators $\mathcal{D}$.

Theorem 2.2 *$\mathcal{D}(X)$ is a free $\mathcal{D}$ -algebra generated by X .*

Proof. Let $\mathcal{A}$ be any $\mathcal{D}$ -algebra and $\varphi : X \rightarrow \mathcal{A}$ a map. We define a linear map $\varphi^* : \mathcal{D}(X) \rightarrow \mathcal{A}$ by

$$\varphi^*(D^{\bar{i}_1}(x_1)D^{\bar{i}_2}(x_2) \cdots D^{\bar{i}_k}(x_k)) = D^{\bar{i}_1}(\varphi(x_1))D^{\bar{i}_2}(\varphi(x_2)) \cdots D^{\bar{i}_k}(\varphi(x_k)).$$

Then, it is easy to check that φ^* is the unique $\mathcal{D}$ -homomorphism such that $\varphi^*\iota = \varphi$, where $\iota : X \rightarrow \mathcal{D}(X)$ is the inclusion map and the proof is complete. $\square$

3 Composition-Diamond lemma for free differential algebras

Let $\mathcal{D} = \{D_j | j \in J\}$, X and J well ordered sets, $D^{\bar{i}}(x) = D_{i_1}D_{i_2} \cdots D_{i_m}(x) \in D^\omega(X)$ and

$$wt(D^{\bar{i}}(x)) = (x; m, i_1, i_2, \dots, i_m).$$

We order $D^\omega(X)$ as follows:

$$D^{\bar{i}}(x) > D^{\bar{j}}(y) \iff wt(D^{\bar{i}}(x)) > wt(D^{\bar{j}}(y)) \text{ lexicographically.}$$

It is easy to check that this order is a well ordering on $D^\omega(X)$.

Now, we order $T = (D^\omega(X))^*$ by deg-lex order $>$, i.e., two words are first compared by length and then lexicographically. We will use this order in the sequel.

Then, for any $0 \neq f \in \mathcal{D}(X)$, we can write

$$f = \alpha_{\bar{f}}\bar{f} + \sum_{i=1}^k \alpha_i u_i,$$

where $\bar{f}, u_i \in T$, $0 \neq \alpha_{\bar{f}}, \alpha_i \in \mathbb{k}$, and $\bar{f} > u_i$. $\bar{f}$ is called the leading term of f and $\alpha_{\bar{f}}$ the leading coefficient. If $\alpha_{\bar{f}} = 1$, we say that f is monic.

The proofs of Lemmas 3.1, 3.2 and 3.3 are straightforward.

Lemma 3.1 *If $u > v$, $u, v \in T$, then for any $a, b \in T$, $aub > avb$. $\square$*

Lemma 3.2 *Let $D^{\bar{i}}(x) = D_{i_1}D_{i_2} \cdots D_{i_m}(x) \in D^\omega(X)$, $u = D^{\bar{i}}(x)v$, $v \in T$. Then, for any $D_j \in \mathcal{D}$, we have*

$$\overline{D_j(u)} = D_j D_{i_1} D_{i_2} \cdots D_{i_m}(x)v. \quad \square$$

Lemma 3.3 *For any $u, v \in T \setminus \{1\}$, if $u > v$, then $\overline{D(u)} > \overline{D(v)}$ for any $D \in \mathcal{D}$. $\square$*

By the above lemmas, the order $>$ on T is monomial in the sense that for any $u, v \in T$,

$$u > v \Rightarrow aub > avb \text{ and } \overline{D(u)} > \overline{D(v)} \text{ for any } a, b \in T, D \in \mathcal{D},$$

where we assume that $v \neq 1$ for the latter case.

Corollary 3.4 *Let $D^{\bar{i}}(x) = D_{i_1}D_{i_2} \cdots D_{i_m}(x) \in D^w(X)$, $u = D^{\bar{i}}(x)v$, $v \in T$. Then, for $D^{\bar{j}} = D_{j_1}D_{j_2} \cdots D_{j_n}$,*

$$\overline{D^{\bar{j}}(u)} = D_{j_1}D_{j_2} \cdots D_{j_n}D_{i_1}D_{i_2} \cdots D_{i_m}(x)v. \quad \square$$

Let the notation be as in Corollary 3.4. For convenience, we denote $\overline{D^{\bar{j}}(u)}$ by $d^{\bar{j}}(u)$.

Definition 3.5 *Let $f, g \in \mathcal{D}(X)$ be monic polynomials and $w, a, b \in T$. Then there are two kinds of compositions.*

(i) *There are two sorts of composition of inclusion:*

If $w = \bar{f} = a \cdot d^{\bar{j}}(\bar{g}) \cdot b$, then the composition is

$$(f, g)_w = f - a \cdot D^{\bar{j}}(g) \cdot b.$$

If $w = d^{\bar{i}}(\bar{f}) = \bar{g} \cdot b$, then the composition is

$$(f, g)_w = D^{\bar{i}}(f) - g \cdot b.$$

(ii) *Composition of intersection:*

If $w = \bar{f} \cdot b = a \cdot d^{\bar{j}}(\bar{g})$ such that $|\bar{f}| + |\bar{g}| > |w|$, then the composition is

$$(f, g)_w = f \cdot b - a \cdot D^{\bar{j}}(g).$$

In this case, we assume that $a, b \neq 1$.

By Lemmas 3.1 and 3.3, we have

Lemma 3.6 *For any composition $(f, g)_w$, we have $\overline{(f, g)_w} < w$. $\square$*

Let $S \subset \mathcal{D}(X)$ be a subset of monic polynomials. Then we define $(S, \mathcal{D})$ -words u_s inductively: for any $s \in S$,

(i) for any $D^{\bar{j}} = D_{j_1}D_{j_2} \cdots D_{j_m}$, $u_s = D^{\bar{j}}(s)$ is an $(S, \mathcal{D})$ -word of $(S, \mathcal{D})$ -length 1;

(ii) if u_s is an $(S, \mathcal{D})$ -word of $(S, \mathcal{D})$ -length k and v is a word in T of length l , then

$$u_s \cdot v \quad \text{and} \quad v \cdot u_s$$

are $(S, \mathcal{D})$ -words of $(S, \mathcal{D})$ -length $k + l$.

Then, any element of $Id(S)$, the $\mathcal{D}$ -ideal of $\mathcal{D}(X)$ generated by S , is a linear combination of $(S, \mathcal{D})$ -words.

By Lemmas 3.1, 3.3 and 3.4, we obtain

Lemma 3.7 *For any $(S, \mathcal{D})$ -word $u_s = a \cdot D^{\bar{j}}(s) \cdot b$, $a, b \in T$, $s \in S$, we have $\overline{u_s} = a \cdot d^{\bar{j}}(\bar{s}) \cdot b$. $\square$*

Definition 3.8 Let $S \subset \mathcal{D}(X)$ be a monic subset and $w \in T$. Then a polynomial $f \in \mathcal{D}(X)$ is said to be trivial modulo (S, w) , denoted by $f \equiv 0 \pmod{(S, w)}$, if

$$f = \sum_i \alpha_i u_{s_i},$$

where each u_{s_i} is an $(S, \mathcal{D})$ -word, $\alpha_i \in \mathbb{k}$ and $\overline{u_{s_i}} < w$.

S is called a Gröbner-Shirshov basis in $\mathcal{D}(X)$ if all compositions of elements of S are trivial modulo S and corresponding w .

Lemma 3.9 S is a Gröbner-Shirshov basis in $\mathcal{D}(X)$ if and only if for any $f, g \in S$,

(i) if $w = d^{\bar{i}}(\bar{f}) = a \cdot d^{\bar{j}}(\bar{g}) \cdot b$ for some $w, a, b \in T$, then

$$D^{\bar{i}}(f) - a \cdot D^{\bar{j}}(g) \cdot b \equiv 0 \pmod{(S, w)};$$

(ii) if $w = d^{\bar{i}}(\bar{f}) \cdot b = a \cdot d^{\bar{j}}(\bar{g})$ for some $w, a, b \in T, a, b \neq 1$ such that $|\bar{f}| + |\bar{g}| > |w|$, then

$$D^{\bar{i}}(f) \cdot b - a \cdot D^{\bar{j}}(g) \equiv 0 \pmod{(S, w)}.$$

Proof. The sufficiency is clear. We only need to prove the necessity.

Firstly, $w = d^{\bar{i}}(\bar{f}) = a \cdot d^{\bar{j}}(\bar{g}) \cdot b$ for some $w, a, b \in T$.

Case 1.1 If $a \neq 1$, then $\exists a' \in T$, such that $d^{\bar{i}}a' = a$. We have $w' = \bar{f} = a'd^{\bar{j}}(\bar{g}) \cdot b$, $(f, g)_{w'} = f - a' \cdot D^{\bar{j}}(g) \cdot b \equiv 0 \pmod{(S, w')}$ and $w = d^{\bar{i}}w'$. Therefore,

$$\begin{aligned} & D^{\bar{i}}(f) - a \cdot D^{\bar{j}}(g) \cdot b \\ &= D^{\bar{i}}(f - a' \cdot D^{\bar{j}}(g) \cdot b) + D^{\bar{i}}(a' \cdot D^{\bar{j}}(g) \cdot b) - a \cdot D^{\bar{j}}(g) \cdot b \\ &= D^{\bar{i}}(f - a' \cdot D^{\bar{j}}(g) \cdot b) + D^{\bar{i}}(a') \cdot D^{\bar{j}}(g) \cdot b + a' \cdot D^{\bar{i}}(D^{\bar{j}}(g) \cdot b) - a \cdot D^{\bar{j}}(g) \cdot b \\ &= D^{\bar{i}}(f - a' \cdot D^{\bar{j}}(g) \cdot b) + d^{\bar{i}}(a') \cdot D^{\bar{j}}(g) \cdot b + \sum_i c_i \cdot D^{\bar{j}}(g) \cdot b + a' \cdot D^{\bar{i}}(D^{\bar{j}}(g) \cdot b) - a \cdot D^{\bar{j}}(g) \cdot b \\ &= D^{\bar{i}}((f, g)_{w'}) + \sum_k c_k \cdot D^{\bar{j}}(g) \cdot b + a' \cdot D^{\bar{i}}(D^{\bar{j}}(g) \cdot b) \end{aligned}$$

where $D^{\bar{i}}(a') = d^{\bar{i}}(a) + \sum_k c_k = a + \sum_k c_k$, $c_k < a$ and all $c_k \cdot D^{\bar{j}}(g) \cdot b$, $a' \cdot D^{\bar{i}}(D^{\bar{j}}(g) \cdot b)$ are $(S, \mathcal{D})$ -words with the leading terms less than w . So we get

$$D^{\bar{i}}(f) - a \cdot D^{\bar{j}}(g) \cdot b \equiv 0 \pmod{(S, w)}.$$

The following three cases can be proved similarly to the Case 1.1 and we omit the details.

Case 1.2 $a = 1$ and $\exists \bar{l}$ such that $\bar{f} = d^{\bar{l}}(\bar{g})b$, $d^{\bar{j}}(\bar{g}) = d^{\bar{i}}(d^{\bar{l}}(\bar{g}))$. Then $w' = \bar{f} = d^{\bar{l}}(\bar{g}) \cdot b$, $w = d^{\bar{i}}w'$ and $(f, g)_{w'} = f - D^{\bar{l}}(g) \cdot b \equiv 0 \pmod{(S, w')}$.

Case 1.3 $a = 1$ and $\exists \bar{l}$ such that $d^{\bar{l}}(\bar{f}) = \bar{g}b$, $d^{\bar{j}}(d^{\bar{l}}(\bar{f})) = d^{\bar{i}}(\bar{f})$. Then $w' = d^{\bar{l}}(\bar{f}) = \bar{g} \cdot b$, $w = d^{\bar{j}}w'$ and $(f, g)_{w'} = D^{\bar{l}}(f) - gb \equiv 0 \pmod{(S, w')}$.

Secondly, $w = d^{\bar{i}}(\bar{f}) \cdot b = a \cdot d^{\bar{j}}(\bar{g})$ for some $w, a, b \in T, a, b \neq 1$ such that $|\bar{f}| + |\bar{g}| > |w|$. Then there exists an $a' \in T$, $a' \neq 1$ such that $a = d^{\bar{i}}(a')$, $w' = \bar{f} \cdot b = a' \cdot d^{\bar{j}}(\bar{g})$, $w = d^{\bar{i}}w'$ and $(f, g)_{w'} = f \cdot b - a' \cdot D^{\bar{j}}(g) \equiv 0 \pmod{(S, w')}$. $\square$

Lemma 3.10 *Let S be a Gröbner-Shirshov basis in $\mathcal{D}(X)$ and $s_1, s_2 \in S$. If*

$$w = a_1 \cdot d^{\bar{i}}(\bar{s}_1) \cdot b_1 = a_2 \cdot d^{\bar{j}}(\bar{s}_2) \cdot b_2$$

for some $a_1, a_2, b_1, b_2 \in T$, then

$$f = a_1 \cdot D^{\bar{i}}(s_1) \cdot b_1 - a_2 \cdot D^{\bar{j}}(s_2) \cdot b_2 \equiv 0 \pmod{(S, w)}.$$

Proof. There are three cases to consider.

Case 1. $d^{\bar{i}}(\bar{s}_1)$ and $d^{\bar{j}}(\bar{s}_2)$ are mutually disjoint. We may assume that $d^{\bar{i}}(\bar{s}_1)$ is at the left of $d^{\bar{j}}(\bar{s}_2)$, i.e.,

$$a_2 = a_1 \cdot d^{\bar{i}}(\bar{s}_1) \cdot a, \quad b_1 = a \cdot d^{\bar{j}}(\bar{s}_2) \cdot b_2$$

where $a \in T$. Then

$$\begin{aligned} f &= a_1 \cdot D^{\bar{i}}(s_1) \cdot b_1 - a_2 \cdot D^{\bar{j}}(s_2) \cdot b_2 \\ &= a_1 \cdot D^{\bar{i}}(s_1) \cdot a \cdot d^{\bar{j}}(\bar{s}_2) \cdot b_2 - a_1 \cdot d^{\bar{i}}(\bar{s}_1) \cdot a \cdot D^{\bar{j}}(s_2) \cdot b_2 \\ &= a_1 \cdot (D^{\bar{i}}(s_1) - d^{\bar{i}}(\bar{s}_1)) \cdot a \cdot D^{\bar{j}}(s_2) \cdot b_2 - a_1 \cdot D^{\bar{i}}(s_1) \cdot a (D^{\bar{j}}(s_2) - d^{\bar{j}}(\bar{s}_2)) \cdot b_2 \\ &= \sum_l \alpha_l a'_l \cdot D^{\bar{j}}(s_2) \cdot b_2 - \sum_k \beta_k a_1 \cdot D^{\bar{i}}(s_1) \cdot b'_k \end{aligned}$$

where $\alpha_l, \beta_k \in \mathbb{k}$, $a'_l, b'_k \in T$, $a'_l < a_1 \cdot d^{\bar{i}}(\bar{s}_1) \cdot a = a_2$, $b'_k < a \cdot d^{\bar{j}}(\bar{s}_2) \cdot b_2 = b_1$. Thus, $f \equiv 0 \pmod{(S, w)}$.

Case 2. One of $d^{\bar{i}}(\bar{s}_1)$ and $d^{\bar{j}}(\bar{s}_2)$ is a subword of the other, say, $w' = d^{\bar{i}}(\bar{s}_1) = a \cdot d^{\bar{j}}(\bar{s}_2) \cdot b$. Then $a_2 = a_1 a$, $b_2 = b b_1$ and

$$\begin{aligned} f &= a_1 \cdot D^{\bar{i}}(s_1) \cdot b_1 - a_2 \cdot D^{\bar{j}}(s_2) \cdot b_2 \\ &= a_1 \cdot D^{\bar{i}}(s_1) \cdot b_1 - a_1 a \cdot D^{\bar{j}}(s_2) \cdot b b_1 \\ &= a_1 \cdot (D^{\bar{i}}(s_1) - a \cdot D^{\bar{j}}(s_2) \cdot b) \cdot b_1 \\ &\equiv 0 \pmod{(S, w)} \end{aligned}$$

since $w = a_1 w' b_1$ and $D^{\bar{i}}(s_1) - a \cdot D^{\bar{j}}(s_2) \cdot b \equiv 0 \pmod{(S, w')}$ by Lemma 3.9.

Case 3. $d^{\bar{i}}(\bar{s}_1)$ and $d^{\bar{j}}(\bar{s}_2)$ have a nonempty intersection as a subword of w , but $d^{\bar{i}}(\bar{s}_1)$ and $d^{\bar{j}}(\bar{s}_2)$ are not subwords of each other. We may assume that

$$a_2 = a_1 a, \quad b_1 = b b_2, \quad w' = d^{\bar{i}}(\bar{s}_1) \cdot b = a \cdot d^{\bar{j}}(\bar{s}_2).$$

Then, we have

$$\begin{aligned} f &= a_1 \cdot D^{\bar{i}}(s_1) \cdot b_1 - a_2 \cdot D^{\bar{j}}(s_2) \cdot b_2 \\ &= a_1 \cdot D^{\bar{i}}(s_1) \cdot b b_2 - a_1 a \cdot D^{\bar{j}}(s_2) \cdot b_2 \\ &= a_1 \cdot (D^{\bar{i}}(s_1) \cdot b - a \cdot D^{\bar{j}}(s_2)) \cdot b_2 \\ &\equiv 0 \pmod{(S, w)} \end{aligned}$$

since $w = a_1 w' b_2$ and also by Lemma 3.9, $D^{\bar{i}}(s_1) \cdot b - a \cdot D^{\bar{j}}(s_2) \equiv 0 \pmod{(S, w')}$.

This completes the proof. $\square$

Lemma 3.11 *Let $S \subset \mathcal{D}(X)$ be a monic subset and $\text{Irr}(S) = \{u \in T \mid u \neq a \cdot d^{\bar{i}}(\bar{s}) \cdot b \text{ for all } s \in S, a, b \in T, \bar{i} \in J^m, m \in \mathbb{N}\}$. Then for any $f \in \mathcal{D}(X)$,*

$$f = \sum_{u_i \leq \bar{f}} \alpha_i u_i + \sum_{\overline{v_{s_j}} \leq \bar{f}} \beta_j v_{s_j}$$

where each $\alpha_i, \beta_j \in \mathbb{k}$, $u_i \in \text{Irr}(S)$ and v_{s_j} is an $(S, \mathcal{D})$ -word.

Proof. Let $f = \sum_i \alpha_i u_i \in \mathcal{D}(X)$, where $0 \neq \alpha_i \in \mathbb{k}$ and $u_1 > u_2 > \dots$. If $u_1 \in \text{Irr}(S)$, then let $f_1 = f - \alpha_1 u_1$. If $u_1 \notin \text{Irr}(S)$, then there exist some $s_1 \in S$, $\bar{i}_1$ and $a_1, b_1 \in T$, such that $\bar{f} = u_1 = a_1 d^{\bar{i}_1}(\bar{s}_1) b_1$. Let $f_1 = f - \alpha_1 a_1 D^{\bar{i}_1}(s_1) b_1$. In both cases, we have $\bar{f}_1 < \bar{f}$. Then the result follows by induction on $\bar{f}$. $\square$

The following theorem is an analogue of the Shirshov's Composition lemma for Lie algebras [25], which was specialized to associative algebras by Bokut [2], see also Bergman [1].

Theorem 3.12 (Composition-Diamond lemma for differential algebras) *Let $\mathcal{D}(X)$ be the free differential algebra with differential operators $\mathcal{D} = \{D_j \mid j \in J\}$, $S \subset \mathcal{D}(X)$ a monic subset, $\text{Id}(S)$ the $\mathcal{D}$ -ideal of $\mathcal{D}(X)$ generated by S and $<$ the order on $T = (D^\omega(X))^*$ as before. Then the following statements are equivalent:*

- (i) S is a Gröbner-Shirshov basis in $\mathcal{D}(X)$.
- (ii) $f \in \text{Id}(S) \Rightarrow \bar{f} = a \cdot d^{\bar{i}}(\bar{s}) \cdot b$ for some $s \in S$, $\bar{i} \in J^m$, $m \in \mathbb{N}$ and $a, b \in T$.
- (iii) $\text{Irr}(S) = \{u \in T \mid u \neq a \cdot d^{\bar{i}}(\bar{s}) \cdot b \text{ for all } s \in S, a, b \in T, \bar{i} \in J^m, m \in \mathbb{N}\}$ is a $\mathbb{k}$ -linear basis of $\mathcal{D}(X|S) = \mathcal{D}(X)/\text{Id}(S)$.

Proof. (i) $\Rightarrow$ (ii). Let S be a Gröbner-Shirshov basis and $0 \neq f \in \text{Id}(S)$. Then

$$f = \sum_{i=1}^k \alpha_i a_i \cdot D^{\bar{j}_i}(s_i) \cdot b_i,$$

where $\alpha_i \in \mathbb{k}$, $a_i, b_i \in T$, $s_i \in S$. Let $w_i = a_i \cdot d^{\bar{j}_i}(\bar{s}_i) \cdot b_i$. We may assume without loss of generality that

$$w_1 = w_2 = \dots = w_l > w_{l+1} \geq w_{l+2} \geq \dots \geq w_k$$

for some $l \geq 1$. We proceed by induction on w_1 and l .

If $l = 1$, then $\bar{f} = w_1 = a_1 \cdot d^{\bar{j}_1}(\bar{s}_1) \cdot b_1$ and we are done.

If $\alpha_1 + \alpha_2 \neq 0$ or $l > 2$, we have $a_1 \cdot d^{\bar{j}_1}(\bar{s}_1) \cdot b_1 = w_1 = w_2 = a_2 \cdot d^{\bar{j}_2}(\bar{s}_2) \cdot b_2$ and furthermore,

$$\begin{aligned} & \alpha_1 a_1 \cdot D^{\bar{j}_1}(s_1) \cdot b_1 + \alpha_2 a_2 \cdot D^{\bar{j}_2}(s_2) \cdot b_2 \\ &= (\alpha_1 + \alpha_2) a_1 \cdot D^{\bar{j}_1}(s_1) \cdot b_1 + \alpha_2 (a_2 \cdot D^{\bar{j}_2}(s_2) \cdot b_2 - a_1 \cdot D^{\bar{j}_1}(s_1) \cdot b_1). \end{aligned}$$

By Lemma 3.10, the last item of the above is a linear combination of $(S, \mathcal{D})$ -words less than w_1 . The result follows by induction on l .

For the case of $\alpha_1 + \alpha_2 = 0$ and $l = 2$, we use induction on w_1 and the result follows.

(ii) $\Rightarrow$ (iii). For any $f \in \mathcal{D}(X)$, by Lemma 3.11, we have

$$f = \sum_{u_i \leq \bar{f}} \alpha_i u_i + \sum_{\overline{v_{s_j}} \leq \bar{f}} \beta_j v_{s_j},$$

where each $\alpha_i, \beta_j \in \mathbb{k}$, $u_i \in \text{Irr}(S)$ and v_{s_j} is an $(S, \mathcal{D})$ -word. Therefore

$$f + \text{Id}(S) = \sum_i \alpha_i (u_i + \text{Id}(S)).$$

On the other hand, suppose that $\sum_i \alpha_i u_i = 0$ in $\mathcal{D}(X)/\text{Id}(S)$, where $\alpha_i \in \mathbb{k}$, $u_i \in \text{Irr}(S)$. This means that $\sum_i \alpha_i u_i \in \text{Id}(S)$ in $\mathcal{D}(X)$. Then all α_i must be equal to zero. Otherwise, $\overline{\sum_i \alpha_i u_i} = u_j \in \text{Irr}(S)$ for some j which contradicts (ii).

(iii) $\Rightarrow$ (i). For any $f, g \in S$, by Lemmas 3.6 and 3.11, we have

$$(f, g)_w = \sum_{u_i \in \text{Irr}(S), u_i < w} \alpha_i u_i + \sum_{\overline{v_{s_j}} < w} \beta_j v_{s_j}.$$

Since $(f, g)_w \in \text{Id}(S)$ and by (iii), we have

$$(f, g)_w = \sum_{\overline{v_{s_j}} < w} \beta_j v_{s_j}.$$

Therefore, S is a Gröbner-Shirshov basis. $\square$

4 Gröbner-Shirshov bases for free Lie-differential algebras

In this section, we define Lie- (commutative-) differential algebras. As applications of the Composition-Diamond lemma for differential algebras (Theorem 3.12), we give Gröbner-Shirshov bases for free Lie-differential algebras and free commutative-differential algebras, respectively.

It is well-known that for an arbitrary algebra $\mathcal{A}$ the set $\text{Der}\mathcal{A}$ is a Lie algebra with respect to the ordinary commutator of linear maps.

Definition 4.1 *Let $(\mathcal{D}, [,])$ be a Lie algebra over $\mathbb{k}$. A Lie-differential algebra (Lie- $\mathcal{D}$ algebra for short) is a $\mathcal{D}$ -algebra $(\mathcal{A}, \partial)$, where $\partial : \mathcal{D} \rightarrow \text{Der}\mathcal{A}$ is a homomorphism of Lie algebras.*

Definition 4.2 *Let X be a set, $(\mathcal{D}, [,])$ a Lie algebra, $L\mathcal{D}(X)$ a Lie- $\mathcal{D}$ algebra and $\iota : X \rightarrow L\mathcal{D}(X)$ an inclusion map. If for any Lie- $\mathcal{D}$ algebra $\mathcal{A}$ and any map $\varphi : X \rightarrow \mathcal{A}$, there exists a unique $\mathcal{D}$ -homomorphism $\varphi^* : L\mathcal{D}(X) \rightarrow \mathcal{A}$ such that $\varphi^* \iota = \varphi$, then $L\mathcal{D}(X)$ is called a free Lie- $\mathcal{D}$ algebra generated by X .*

The proof of the following theorem is straightforward and we omit the details.

Theorem 4.3 *Let $(\mathcal{D}, [\cdot, \cdot])$ be a Lie algebra with a $\mathbb{k}$ -basis $\{D_j \mid j \in J\}$ and $[D_i, D_j] = D_i D_j - D_j D_i = \sum_r \alpha_{ij}^r D_r$, $i, j \in J$, where each $\alpha_{ij}^r \in \mathbb{k}$. Let $\mathcal{D}(X)$ be a free $\mathcal{D}$ -algebra and*

$$S = \{D_i D_j(u) - D_j D_i(u) - \sum_r \alpha_{ij}^r D_r(u) \mid u \in T, i, j \in J\}.$$

Then $\mathcal{D}(X|S) = \mathcal{D}(X)/Id(S)$ is a free Lie- $\mathcal{D}$ algebra generated by X . $\square$

Let

$$\begin{aligned} H &= \{D_{i_1} \cdots D_{i_m}(x) \mid i_1 \leq \cdots \leq i_m, i_1, \dots, i_m \in J, m \in \mathbb{N}, x \in X\}, \\ D^{\bar{i}}(x) &= D_{i_1} \cdots D_{i_m}(x), \\ S_0 &= \{D_p D_q D^{\bar{i}}(x) - D_q D_p D^{\bar{i}}(x) - \sum_r \alpha_{pq}^r D_r D^{\bar{i}}(x) \mid p > q, p, q \in J, D^{\bar{i}}(x) \in H, x \in X\}. \end{aligned}$$

Lemma 4.4 *$Irr(S_0) = H^*$, where H^* is the free monoid generated by H . $\square$*

Lemma 4.5 *Suppose S and S_0 are sets defined as above. Then, they generate the same $\mathcal{D}$ -ideal in $\mathcal{D}(X)$, i.e.,*

$$Id(S) = Id(S_0).$$

Proof. Since S_0 is a subset of S , it suffices to prove that $\mathcal{D}(X|S_0)$ is a Lie- $\mathcal{D}$ algebra. We need only prove that in $\mathcal{D}(X|S_0)$, for any $u \in T$,

$$D_i D_j(u) - D_j D_i(u) - \sum_r \alpha_{ij}^r D_r(u) = 0.$$

By Lemma 3.11, it suffices to prove that for any $v \in Irr(S_0)$,

$$D_i D_j(v) - D_j D_i(v) - \sum_r \alpha_{ij}^r D_r(v) = 0.$$

We use induction on the length of v .

If $|v| = 1$, then the result is trivial.

If $|v| \geq 2$, i.e., $v = D^{\bar{i}}(x)v_1$ where $D^{\bar{i}}(x) \in H$, $1 \neq v_1 \in H^*$, then by induction,

$$\begin{aligned} & D_i D_j(v) - D_j D_i(v) - \sum_r \alpha_{ij}^r D_r(v) \\ &= D_i D_j(D^{\bar{i}}(x)v_1) - D_j D_i(D^{\bar{i}}(x)v_1) - \sum_r \alpha_{ij}^r D_r(D^{\bar{i}}(x)v_1) \\ &= (D_i D_j D^{\bar{i}}(x) - D_j D_i D^{\bar{i}}(x) - \sum_r \alpha_{ij}^r D_r D^{\bar{i}}(x))v_1 \\ &\quad + D^{\bar{i}}(x)(D_i D_j(v_1) - D_j D_i(v_1) - \sum_r \alpha_{ij}^r D_r(v_1)) \\ &= 0. \quad \square \end{aligned}$$

By Lemma 3.11, we have

Lemma 4.6 For any $p, q \in J$ and $D^{\bar{i}}x \in D^\omega(X)$, if $p > q$, then

$$D_p D_q D^{\bar{i}}(x) - D_q D_p D^{\bar{i}}(x) - \sum_r \alpha_{pq}^r D_r D^{\bar{i}}(x) \equiv 0 \quad \text{mod}(S_0, w)$$

where $w \in D^\omega(X)$ and $w > D_p D_q D^{\bar{i}}x$. $\square$

Theorem 4.7 Let $(\mathcal{D}, [,]) be a Lie algebra with a $\mathbb{k}$ -basis $\{D_j \mid j \in J\}$ and $[D_i, D_j] = D_i D_j - D_j D_i = \sum_r \alpha_{ij}^r D_r$, $i, j \in J$, where each $\alpha_{ij}^r \in \mathbb{k}$. Let $\mathcal{D}(X)$ be the free differential algebra with differential operators $\mathcal{D}$. Let the order $<$ on T be as before and$

$$S_0 = \{D_p D_q D_{i_1} \cdots D_{i_m}(x) - D_q D_p D_{i_1} \cdots D_{i_m}(x) - \sum_r \alpha_{pq}^r D_r D_{i_1} \cdots D_{i_m}(x) \mid \\ p > q, i_1 \leq \cdots \leq i_m, p, q, i_1, \dots, i_m \in J, m \geq 0, x \in X\}.$$

Then

- (i) S_0 is a Gröbner-Shirshov basis in $\mathcal{D}(X)$.
- (ii) H^* is a $\mathbb{k}$ -basis of the free Lie- $\mathcal{D}$ algebra $\mathcal{D}(X|S)$ generated by X , where $H = \{D_{i_1} \cdots D_{i_m}(x) \mid i_1 \leq \cdots \leq i_m, i_1, \dots, i_m \in J, m \in \mathbb{N}, x \in X\}$ and H^* is the free monoid generated by H .

Proof. (i). The possible composition is only the case of inclusion $(f, g)_w = f - D_p g$ where

$$f = D_p D_q D_{i_1} \cdots D_{i_m}(x) - D_q D_p D_{i_1} \cdots D_{i_m}(x) - \sum_r \alpha_{pq}^r D_r D_{i_1} \cdots D_{i_m}(x), \\ g = D_q D_{i_1} D_{i_2} \cdots D_{i_m}(x) - D_{i_1} D_q D_{i_2} \cdots D_{i_m}(x) - \sum_r \alpha_{q i_1}^r D_r D_{i_2} \cdots D_{i_m}(x), \\ w = \bar{f} = \overline{D_p g}, p > q > i_1, i_1 \leq i_2 \leq \cdots \leq i_m, m \geq 1.$$

Now, we prove that $(f, g)_w \equiv 0 \text{ mod}(S_0, w)$. Denote by $u = D_{i_2} \cdots D_{i_m}(x)$. Then

$$(f, g)_w = D_p D_{i_1} D_q u + \sum_r \alpha_{q i_1}^r D_p D_r u - D_q D_p D_{i_1} u - \sum_r \alpha_{pq}^r D_r D_{i_1} u.$$

Let $A = D_p D_{i_1} D_q u$, $B = -D_q D_p D_{i_1} u$ and $C = \sum_r \alpha_{q i_1}^r D_p D_r u - \sum_r \alpha_{pq}^r D_r D_{i_1} u$. Then, by Lemma 4.6, we have

$$A = D_p D_{i_1} D_q u \\ \equiv D_{i_1} D_p D_q u + \sum_r \alpha_{p i_1}^r D_r D_q u \\ \equiv D_{i_1} D_q D_p u + \sum_r \alpha_{pq}^r D_{i_1} D_r u + \sum_r \alpha_{p i_1}^r D_r D_q u \quad \text{mod}(S_0, w)$$

and

$$B = -D_q D_p D_{i_1} u \\ \equiv -D_q D_{i_1} D_p u - \sum_r \alpha_{p i_1}^r D_q D_r u \\ \equiv -D_{i_1} D_q D_p u - \sum_r \alpha_{q i_1}^r D_r D_p u - \sum_r \alpha_{p i_1}^r D_q D_r u \quad \text{mod}(S_0, w).$$

Since $(\mathcal{D}, [,]) is a Lie algebra, we have$

$$[D_i, D_j] = -[D_j, D_i] \text{ and } [[D_i, D_j], D_k] + [[D_k, D_i], D_j] + [[D_j, D_k], D_i] = 0.$$

It follows that

$$\alpha_{ij}^s = -\alpha_{ji}^s \text{ and } \sum_{s,t} \alpha_{ij}^s \alpha_{sk}^t + \sum_{s,t} \alpha_{ki}^s \alpha_{sj}^t + \sum_{s,t} \alpha_{jk}^s \alpha_{si}^t = 0.$$

Thus,

$$\begin{aligned} (f, g)_w &= A + B + C \\ &\equiv \sum_r \alpha_{pq}^r D_{i_1} D_r u + \sum_r \alpha_{pi_1}^r D_r D_q u + \sum_r \alpha_{qi_1}^r D_p D_r u \\ &\quad - \sum_r \alpha_{qi_1}^r D_r D_p u - \sum_r \alpha_{pi_1}^r D_q D_r u - \sum_r \alpha_{pq}^r D_r D_{i_1} u \\ &\equiv \sum_{r,l} \alpha_{pq}^r \alpha_{i_1 r}^l D_l u + \sum_{r,l} \alpha_{pi_1}^r \alpha_{rq}^l D_l u + \sum_{r,l} \alpha_{qi_1}^r \alpha_{pr}^l D_l u \\ &\equiv \left(\sum_{r,l} \alpha_{pq}^r \alpha_{i_1 r}^l + \sum_{r,l} \alpha_{pi_1}^r \alpha_{rq}^l + \sum_{r,l} \alpha_{qi_1}^r \alpha_{pr}^l \right) D_l u \\ &\equiv 0 \pmod{(S_0, w)}. \end{aligned}$$

This shows (i).

(ii) follows from Theorem 3.12, Theorem 4.3, Lemma 4.4 and Lemma 4.5. $\square$

Remark: Let $\mathcal{A}$ be a differential algebra with differential operators $\mathcal{D}$. If $\mathcal{D}$ is a commutative $\mathbb{k}$ -algebra, then $\mathcal{A}$ is called a commutative-differential algebra. Clearly, a commutative-differential algebra is a special Lie-differential algebra. Therefore, all results in this section are true for commutative-differential algebras.

References

- [1] G. M. Bergman, “The Diamond Lemma for Ring Theory”, *Adv. in Math.*, **29**(1978), p.178.
- [2] L. A. Bokut, “Imbeddings into Simple Associative Algebras”, *Algebra i Logika*, **15**(1976), p.117.
- [3] L. A. Bokut, Yuqun Chen, “Gröbner-Shirshov Bases: some new Results”, *Proceedings of the Second International Congress in Algebra and Combinatorics*, World Scientific, 2008, p.35.
- [4] L. A. Bokut, Yuqun Chen and Yongshan Chen, “Composition-Diamond Lemma for Tensor Product of Free Algebras”, *J. Algebra*, to appear. [arXiv.org/abs/0804.2115](https://arxiv.org/abs/0804.2115).
- [5] L. A. Bokut, Yuqun Chen and Cihua Liu, “Gröbner-Shirshov Bases for Dialgebras”, [arXiv.org/abs/0804.0638](https://arxiv.org/abs/0804.0638).

- [6] L. A. Bokut, Yuqun Chen and Jianjun Qiu, “Gröbner-Shirshov Bases for Associative Algebras with Multiple Operators and Free Rota-Baxter Algebras”, arXiv.org/abs/0805.0640.
- [7] L. A. Bokut, Y. Fong and W. F. Ke, “Composition Diamond Lemma for Associative Conformal Algebras”, *J. Algebra*, **272**(2004), p.739.
- [8] L. A. Bokut and K. P. Shum, “Relative Gröbner-Shirshov Bases for Algebras and Groups”, *Algebra and Analysis*, **19**(2007), p.1. (in Russian)
- [9] B. Buchberger, “An Algorithm for Finding a Basis for the Residue Class Ring of a Zero-dimensional Polynomial Ideal”, Ph.D. thesis, University of Innsbruck, Austria, (1965). (in German)
- [10] B. Buchberger, “An Algorithmical Criteria for the Solvability of Algebraic Systems of Equations”, *Aequationes Math.*, **4**(1970), p.374. (in German)
- [11] E. S. Chibrikov, “On Free Lie Conformal Algebras”, *Vestnik Novosibirsk State University*, **4**(2004), p.65.
- [12] Yuqun Chen, Yongshan Chen and Chanyan Zhong, “Composition-Diamond Lemma for Modules”, *Czech. J. Math.*, to appear. arXiv.org/abs/0804.0917.
- [13] D. R. Farkas, C. D. Feustel and E. L. Green, “Synergy in the Theories of Gröbner Bases and Path Algebras”, *Can. J. Math.*, **45**(1993), p.727.
- [14] H. Hironaka, “Resolution of Singulatities of an Algebtaic Variety over a Field if Characteristic Zero, I, II”, *Ann. Math.*, **79**(1964), p.109, p.205.
- [15] L. Hellström, “The Diamond Lemma for Power Series Algebras”, Ph.D. thesis, 2002.
- [16] S.-J. Kang and K.-H. Lee, “Gröbner-Shirshov Bases for Representation Theory”, *J. Korean Math. Soc.*, **37**(2000), p.55.
- [17] S.-J. Kang and K.-H. Lee, “Linear Algebraic Approach to Gröbner-Shirshov Bases Theory”, *J. Algebra*, **313**(2007), p.988.
- [18] Y. Kobayashi, “Gröbner Bases on Algebras Based on Well-ordered Semigroups”, *Math. Appl. Sci. Tech.*, to appear.
- [19] A. A. Mikhalev, “Shirshov’s Composition Techniques in Lie Superalgebras (Non-commutative Gröbner Bases)”, *Trudy Sem. Petrovsk*, **18**(1995), p.277 (in Russian). English translation: *J. Math. Sci.*, **80**(1996), p.2153.
- [20] A. A. Mikhalev and E. A. Vasilieva, “Standard Bases of Ideals of Free Supercommutative Polynomail Algebra (ε -Grobner Bases)”, *Proc. Second International Taiwan-Moscow Algebra Workshop*, Springer-Verlag, 2003.
- [21] A. A. Mikhalev and A. A. Zolotykh, “Standard Gröbner-Shirshov Bases of Free Algebras over Rings, I. Free Associative Algebras”, *International Journal of Algebra and Computation*, **8**(1998), p.689.

- [22] M. H. A. Newman, “On Theories with a Combinatorial Definition of ‘Equivalence’”, *Ann. of Math.*, **43**(1942), p.223.
- [23] A. I. Shirshov, “On Free Lie Rings”, *Mat. Sb.*, **45(87)**(1958), p.113. (in Russian)
- [24] A. I. Shirshov, “Some Algorithmic Problem for ε -Algebras”, *Sibirsk. Mat. Z.*, **3**(1962), p.132. (in Russian)
- [25] A. I. Shirshov, “Some Algorithmic Problem for Lie Algebras”, *Sibirsk. Mat. Z.*, **3**(1962), p.292 (in Russian). English translation: *SIGSAM Bull.*, **33(2)**(1999), p.3.
- [26] T. Stokes, “Gröbner-Shirshov Bases in Exterior Algebras”, *J. Automated Reasoning*, **6**(1990), p.233.
- [27] A. I. Zhukov, “Complete Systems of Defining Relations in Non-associative Algebras”, *Mat. Sbornik*, **69(27)**(1950), p.267.